

Diabetes-induced glomerular dysfunction: Links to a more reduced cytosolic ratio of NADH/NAD⁺

RONALD G. TILTON, LISA D. BAIER, JUDITH E. HARLOW, SAMUEL R. SMITH, EVA OSTROW, and JOSEPH R. WILLIAMSON

Department of Pathology, Washington University School of Medicine, Saint Louis, Missouri, USA

Diabetes-induced glomerular dysfunction: Links to a more reduced cytosolic ratio of NADH/NAD⁺. These studies were undertaken to examine effects of elevated glucose levels on glycolysis, sorbitol pathway activity, and the cytosolic redox state of NADH/NAD⁺ in isolated glomeruli. Blood-free glomeruli were isolated from kidneys of male, Sprague-Dawley rats using standard sieving techniques, then incubated for one hour at 37°C, pH 7.4, pO₂ ~500 torr, in Krebs bicarbonate/Hepes buffer containing 5 or 30 mM glucose. Elevated glucose levels increased glucose 6-phosphate, fructose 6-phosphate, total triose phosphates, lactate, the lactate/pyruvate ratio, sorbitol, and fructose, but did not affect *sn*-glycerol 3-phosphate, pyruvate, or *myo*-inositol levels. The more reduced glomerular cytosolic redox state (manifested by the tissue lactate/pyruvate ratio) induced by 30 mM glucose was completely abrogated by aldose reductase inhibitors added to the diet two to seven days prior to glomerular isolation. These observations, coupled with evidence linking glucose- and diabetes-induced glomerular dysfunction to increased sorbitol pathway metabolism, support the hypothesis that metabolic imbalances associated with a more reduced ratio of cytosolic NADH/NAD⁺ (resulting from increased glucose metabolism via the sorbitol pathway) play an important role in mediating glucose- and diabetes-induced glomerular dysfunction.

Glomerular dysfunction is readily demonstrable in humans and in animals shortly after the onset of poorly controlled diabetes [1–6]. These glomerular changes, manifested as increases in blood flow, GFR, and microalbuminuria, together with associated increases in blood flow and/or vascular albumin permeation in ocular tissues, sciatic nerve, and aorta of diabetic rats, have been linked to increased metabolism of glucose via the sorbitol pathway by many [5–11] but not all [12] investigators. Increased GFR in human diabetics also is attenuated by treatment with an inhibitor of aldose reductase [13]. The precise nature of the metabolic imbalances that mediate sorbitol pathway-linked glomerular and other vascular (and neural) dysfunction, however, remains unclear.

Topical application of glucose or sorbitol to newly formed granulation tissue vessels in skin chambers increases vascular albumin permeation and blood flow in nondiabetic rats to levels observed in diabetic rats, but in the absence of insulin deficiency and associated systemic hormonal and metabolic imbalances characteristic of the diabetic milieu [6, 14]. As shown in Figure 1, these glucose-induced vascular changes are prevented

by coadministration of: 1) pyruvate [15–17] which also prevents associated increases in tissue levels of 1,2-diacyl-*sn*-glycerol (DAG) in extracts of granulation tissue [15]; 2) inhibitors of aldose reductase [14] which also prevent glucose-induced increases in the ratio of lactate/pyruvate [18] (the most reliable parameter of cytosolic NADH/NAD⁺ [19]); 3) staurosporine [15], an inhibitor of protein kinase C; and 4) *myo*-inositol [17]. The mechanism by which *myo*-inositol prevents sorbitol pathway-linked vascular dysfunction remains unclear. Taken together, these observations suggest that glucose-induced vascular dysfunction is linked to elevated tissue levels of DAG and associated activation of protein kinase C [15] resulting from a more reduced cytosolic ratio of NADH/NAD⁺. This redox change results from increased oxidation of sorbitol to fructose which is coupled to reduction of NAD⁺ to NADH (Fig. 1) [6, 11].

Glucose-induced increased *de novo* synthesis of DAG, increased tissue levels of DAG, and associated activation of protein kinase C also have been observed in isolated glomeruli [20, 21] and in cultured bovine retinal capillary endothelial cells [22, 23]; an increase in DAG mass has been reported in whole retina and in myocardium of diabetic rats [22, 24]; and activation of protein kinase C has been observed in bovine aortic smooth muscle cells and endothelial cells exposed to elevated glucose levels in vitro [22]. A more reduced cytosolic ratio of NADH/NAD⁺ could promote increased *de novo* synthesis of DAG through two mechanisms (Fig. 1). First, it would favor reduction of dihydroxyacetone phosphate (DHAP) to *sn*-glycerol 3-phosphate (*sn*G3P), the first step in the pathway for *de novo* synthesis of DAG. Second, it would tend to increase the availability of substrate (DHAP) by impairing oxidation of glyceraldehyde 3-phosphate (GAP) to 1,3-bisphosphoglycerate (1,3 DPG). Evidence of a more reduced cytosolic ratio of NADH/NAD⁺ has been observed in a variety of tissues of diabetic animals as well as in cells and tissues exposed to elevated glucose levels in vitro [6, 18, 19, 25–27].

In light of these considerations, the present studies were undertaken to examine effects on isolated glomeruli of elevated glucose levels in vitro on cytosolic NADH/NAD⁺ reflected by the lactate/pyruvate ratio. The likelihood that acute glucose-induced in vitro metabolic imbalances of this nature may mediate increased GFR in diabetic subjects is supported by evidence that GFR is acutely increased in human subjects and in nondiabetic animals by brief intravascular infusion of glucose




Fig. 1. Interrelated metabolic imbalances linked to glucose- and diabetes-induced vascular dysfunction. *Pharmacologic interventions which prevent glucose-induced vascular dysfunction are indicated by an asterisk and include: aldose reductase inhibitors (ARI), aminoguanidine (AG), staurosporine (stauro), pyruvate, *myo*-inositol (*myo*), diphenhydramine (DPH), and acetyl-L-carnitine (Acetyl-L-C). *sn*G3P = *sn*-glycerol 3-phosphate, FA = free fatty acid, GA = glyceraldehyde, LcAcyl-C = long-chain acylcarnitine, LcAcyl-CoA = long-chain acyl coenzyme A, NEG = nonenzymatic glycation, PKC = protein kinase C, PI = phosphatidylinositol. Substances which mimic glucose effects on vascular function are italicized and underlined. See references [6, 11] for further discussion of the effects of pharmacologic agents depicted in this figure.

[28, 29] or of lactate [30]. Lactate infusion would increase intra- and extracellular lactate/pyruvate ratios resulting in a more reduced cytosolic ratio of NADH/NAD⁺ and hindering oxidation of NADH to NAD⁺ (coupled to reduction of pyruvate to lactate).

Methods

Animal protocols

All rats used in these experiments were housed and cared for in accordance with the guidelines of the University Committee for the Humane Care of Laboratory Animals and in accordance with NIH guidelines on laboratory animal welfare. Rats were housed individually, allowed food and water *ad libitum*, and were on a 12-hour light/dark cycle. Glomeruli were obtained from kidneys of normal male, Sprague-Dawley rats weighing ~250 g.

Glomerular isolation protocols

For each glomerular isolation, five male Sprague-Dawley rats were anesthetized with sodium pentobarbital (35 to 45 mg/kg body weight) and anticoagulated with sodium heparin (2000 units i.p.) 10 to 15 minutes prior to surgery. A midline abdominal incision was made to expose both kidneys as well as the aorta and vena cava distal to the renal arteries. The aorta was occluded proximal to the renal arteries, a 21-gauge needle

(attached to a 30 ml syringe) was inserted into the abdominal aorta below the renal arteries, a cut was made in the vena cava at the level of the renal arteries, and both kidneys were perfused by hand with ~20 ml of Krebs bicarbonate/Hepes (KH) buffer (containing in mM: NaCl, 116; KCl, 5.4; CaCl₂ · 2H₂O, 1.8; NaH₂PO₄ · H₂O, 1.0; MgSO₄ · 7H₂O, 0.8; NaHCO₃, 15.0; glucose, 5.0; and Hepes, 18.0), pH 7.4, 4°C. Both kidneys blanched within seconds after initiation of buffer perfusion and were moved ~20 seconds later and placed in a beaker containing KH buffer on ice. Immediately following the removal of kidneys from five rats, the capsules were stripped from all kidneys, medullary tissue was removed and discarded, and the pooled cortical tissue was minced on a ice-chilled cutting block into a fine paste with a razor blade. Glomeruli were isolated by sequential sieving at room temperature through three stainless steel sieves prewet with the KH buffer as described by Schlondorff et al [31]. The 8-inch diameter sieves were (from top to bottom) a Tyler equivalent 100 mesh (pore size 150 μm), a Tyler equivalent 170 mesh (pore size 90 μm), and a Tyler equivalent 270 mesh (pore size 53 μm). The cortical paste was gently forced through each sieve using the bottom of a 200 ml glass beaker and 2 to 3 liters of the KH buffer at room temperature for each sieve.

Glomeruli were flushed into a 50-ml conical bottom polypropylene tube using KH buffer and centrifuged for one minute at

200 rpm. The supernatant was removed and the glomeruli were resuspended in 10 ml of KH buffer containing 0.05% bovine serum albumin and 75 μM *myo*-inositol (incubation buffer). After gassing with 95% O_2 and 5% CO_2 ($p\text{O}_2$, ~ 500 torr) at 37°C, the glomeruli were allowed to rest for 15 minutes. The glomeruli were then centrifuged (2000 rpm for 30 seconds), the supernatant was removed, and the glomerular pellet was resuspended in fresh incubation buffer in a total volume of ~ 2.6 ml. Six hundred microliter aliquots of well-mixed glomerular suspension (20,000 to 30,000 glomeruli) were added to four 5-ml sialinized vacutainer tubes in a 37°C (shaking) water bath. Five 10- μl samples were removed for glomerular counting and to check their purity by light microscopy. The mean ratio of glomeruli to tubular fragments for all preparations was 15:1 with a standard deviation of 2.6%. After addition of substrates (in a volume of 25 μl), the tubes were gassed with humidified 95% O_2 and 5% CO_2 for five minutes and then tightly capped and incubated for one hour at 37°C.

Experimental protocols

In the first set of preliminary experiments, glomeruli were incubated in 5 or 50 mM glucose for 30, 60, or 120 minutes or in 5, 15, or 30 mM glucose for 60 minutes. Incubations were terminated by rapidly adding 60 μl of 12 N perchloric acid to the incubation buffer with shaking and the extract (tissue + medium) was saved for measurement of glycolytic intermediates. In the second series of experiments, glomeruli were incubated in 5 or 30 mM glucose \pm 3 mM pyruvate or 0.07 mM tolrestat (*N*-[[5-(trifluoromethyl)-6-Methoxy-1-naphthalenyl]-thioxomethyl]-*N* methylglycine) and terminated as described above for assessment of glycolytic intermediates or as described below for quantification of glomerular polyol and *myo*-inositol content. A third set of experiments was performed on glomeruli removed from normal rats fed standard ground chow or ground chow containing tolrestat in an amount to provide 0.2 mmol/kg body weight/day or AL-1576 (spiro-(2,7-difluoro-9H-fluoren-9,4'-imidazolidine)2',5'-dione) in an amount to provide ~ 0.02 mmol/kg body weight/day for two to seven days prior to glomerular isolation. In these experiments, glomeruli from control rats were incubated only in 5 or 30 mM glucose while glomeruli from rats treated with tolrestat or AL-1576 were incubated in the same buffers containing 0.07 mM tolrestat or 0.007 mM AL-1576 added in a volume of 5 μl . After a one hour incubation, glomeruli were extracted for assessment of glycolytic intermediates (tissue + medium). In a fourth experiment, glomeruli were obtained from kidneys of rats fed standard ground chow or ground chow containing tolrestat as described above, then incubated for one hour at 37°C in 2.5 ml KH buffer containing 5 or 30 mM glucose + 0.07 mM tolrestat. Glomeruli were allowed to settle to the bottom of the incubation tube during the last minute of incubation, then rapidly pipetted into a 1.5 ml microfuge tube on ice and centrifuged at 14,000 g for 30 seconds at 4°C. The remaining medium was removed and 50 μl of 12 N perchloric acid was added to the pellet. The extract (tissue only) was saved for determination of glomerular lactate and pyruvate.

Metabolic and biochemical assays

Glycolytic intermediates. Five to ten minutes following addition of perchloric acid as described above, the tubes were

centrifuged, the supernatant was removed, the pH was adjusted to ~ 3.5 , and the supernatant was assayed for fructose, *sn*-glycerol 3-phosphate, and glycolytic metabolites (glucose 6-phosphate, G6P; fructose 6-phosphate, F6P; fructose 1,6-bisphosphate, dihydroxyacetone phosphate, and glyceraldehyde 3-phosphate measured together as total triose phosphates; pyruvate, and lactate) by standard enzymatic methods [32, 33]. The pellet was saved for measurement of hydroxyproline [34] to which all the metabolic data were normalized. Prior to fructose analysis extracts were treated with glucose oxidase [35] to prevent interference with measurement of fructose by high glucose levels.

The effect of elevated glucose levels on the cytosolic ratio of NADH/NAD⁺ was not measured directly, but was inferred from changes in the ratio of glomerular lactate/pyruvate; the cytosolic ratio of these metabolites is a more reliable parameter of the cytosolic ratio of free NADH/NAD⁺ than measurement of the pyridine nucleotides themselves in tissue extracts [19, 36]. Since, depending on the experimental conditions and interventions performed, changes in lactate/pyruvate in the incubation medium at the end of an incubation may not correspond to the ratio of lactate/pyruvate in glomerular cytosol, experiments were performed to verify that changes in lactate/pyruvate in glomeruli plus medium (measured in the first three series of experiments) did indeed reflect the direction of changes in glomerular cytosol.

Glomerular polyol and *myo*-inositol. Following the one hour incubation, glomeruli were immediately centrifuged at 4°C, the pellet was resuspended in 1 ml of 0.02% sodium azide in water, internal standard was added, and the glomeruli were heated at 100°C for 15 minutes. The glomerular extracts were deproteinized with 0.3 N barium hydroxide and 0.3 N zinc sulfate, centrifuged, and sorbitol and *myo*-inositol in the supernatant were quantified (as their butyl boronate derivatives) by gas chromatography/mass spectrometry [37]. The pellet was saved for measurement of hydroxyproline content.

Statistical analysis. All data are expressed as means \pm standard deviations of untransformed data. Since all experiments were performed on paired incubations, overall differences among groups for each parameter were assessed by a repeated measures analysis of variance with the SAS general linear-models procedure [38]. If the repeated measures analysis of variance test indicated that differences among groups were statistically significant at $P < 0.05$ for a given parameter, pairwise comparisons were assessed by least-square means. Because of the large variance in some parameters, a natural log transformation of the data was performed before assessment of differences between groups. The effects of aldose reductase inhibitors given *in vivo* on lactate/pyruvate ratios in glomeruli after incubation in 5 versus 30 mM glucose were assessed using a paired *t*-test.

Results

Effects of elevated glucose levels on glomerular glycolysis

In preliminary experiments (data not shown) glycolytic intermediates were measured in glomeruli incubated in 5 or 50 mM glucose for 30, 60, or 120 minutes. All glycolytic intermediates in the 5 mM glucose incubations reached a steady state level by 60 minutes or remained constant for 120 minutes of incubation,




Fig. 2. Glycolytic intermediates (glucose 6-phosphate, fructose 6-phosphate, total triose phosphates, and *sn*-glycerol 3-phosphate) in glomeruli incubated for one hour at 37°C in 5, 15, or 30 mM glucose. Metabolites, expressed as nmol/μg hydroxyproline, are shown as the mean \pm 1 SD ($N = 7$ for each glucose concentration). Significantly different from 5 mM glucose: * $P < 0.01$, † $P < 0.03$.

with the exception of triose phosphates which increased 1.6-fold in 120 versus 30 minute incubations. Fifty millimolars of glucose increased G6P, F6P, triose phosphates, and lactate (but not pyruvate) after 30 minutes of incubation ($P \leq 0.05$). All of these metabolites remained elevated throughout 120 minutes of incubation except F6P. Pyruvate levels were significantly increased at 60 and 120 minutes ($P < 0.025$). Based on these data, subsequent incubations were carried out for 60 minutes.

Incubation of glomeruli for one hour in medium containing 5, 15, or 30 mM glucose (Figs. 2 and 3) resulted in a concentration-dependent increase in glomerular levels/production of G6P, F6P, total triose phosphates, and lactate ($P < 0.008$, 0.01, 0.02, and 0.03, respectively). Pyruvate and *sn*G3P levels remained constant between 5 and 30 mM glucose and the lactate/pyruvate ratio (Fig. 3C) increased from 11.7 ± 3.3 at 5 mM glucose to 14.8 ± 3.6 at 30 mM glucose ($P < 0.003$). Based on the results of these experiments, all subsequent incubations were performed for 60 minutes in either 5 or 30 mM glucose \pm pharmacologic agents.

Effects of pyruvate and aldose reductase inhibitors added in vitro on glucose-induced metabolic changes

The effects of 30 mM glucose on glomerular levels/production of glycolytic intermediates in these experiments (Figs. 4 and 5) are consistent with those observed in the preliminary studies (Figs. 2 and 3). Sorbitol and fructose levels were increased significantly ($P < 0.001$ and $P < 0.014$, respectively) and *myo*-inositol content decreased slightly (but insignificantly) in glomeruli incubated in 30 versus 5 mM glucose (Fig. 6).

Coadministration of 3 mM pyruvate with 30 mM glucose was associated with a small increase in lactate levels (statistically insignificant vs. 30 mM glucose), an $\sim 70\%$ decrease in triose phosphates ($P < 0.001$ vs. 30 mM glucose) and a similar $\sim 70\%$ decrease in *sn*G3P ($P < 0.0002$ vs. 30 mM glucose), but had no effect on F6P or G6P (Figs. 4 and 5). Pyruvate also decreased glomerular sorbitol levels by $\sim 60\%$ ($P < 0.01$ vs. 30 mM

glucose) but had no effect on fructose or *myo*-inositol levels (Fig. 6).

In vitro addition of 0.07 mM tolrestat to 30 mM glucose had no effect on 30 mM glucose-induced changes in lactate, pyruvate, lactate/pyruvate ratio, or triose phosphates (Figs. 4 and 5). In contrast, tolrestat significantly reduced glomerular levels of *sn*G3P (Fig. 4D). Tolrestat also largely prevented 30 mM glucose-induced increases in glomerular sorbitol levels (Fig. 6A) but had no effect on fructose or *myo*-inositol levels (Fig. 6B and C). While the 30 mM glucose-induced increase in lactate/pyruvate ratio was unaffected by the addition of 0.07 mM tolrestat in vitro (Fig. 5C), the increase was abolished in corresponding experiments with glomeruli from rats given tolrestat or AL-1576 in the diet prior to isolation of glomeruli (Fig. 7A).

Effects of elevated glucose and aldose reductase inhibitors on cytosolic redox state of glomeruli

To obtain a more reliable assessment of glomerular cytosolic NADH/NAD⁺, lactate/pyruvate ratios were measured in glomeruli rapidly separated from incubation medium. As shown in Figure 7B, lactate/pyruvate ratios in 5 mM glucose-incubated glomeruli were almost twice those in extracts of glomeruli plus medium (Fig. 7A). Nevertheless, the ratios increased $\sim 30\%$ after 60 minutes of incubation in 30 mM (31.2 ± 11.6) versus 5 mM (23.8 ± 7.5) glucose, closely corresponding to the increases observed in tissue plus medium after the same duration of incubation (Fig. 7A). The lactate/pyruvate in glomeruli from rats fed ARI for seven days and incubated in 5 mM glucose plus ARI was significantly lower than that in glomeruli from control rats; 30 mM glucose-induced increase in lactate/pyruvate was prevented in 30 mM glucose-incubated glomeruli from ARI-fed rats.

Metabolite levels (in medium plus glomeruli) for glomeruli isolated from ARI-fed versus untreated rats following incubation in 5 and 30 mM glucose are summarized in Figure 8.




Fig. 3. Effects of increasing glucose concentration (5, 15, and 30 mM glucose) on lactate, pyruvate, and lactate/pyruvate ratio, in glomeruli incubated for one hour at 37°C. Metabolites, expressed as nmol/ μ g hydroxyproline, are shown as the mean \pm 1 SD for each glucose concentration ($N = 7$). Significantly different from 5 mM glucose: * $P < 0.003$, † $P < 0.03$, ‡ $P < 0.05$.

Elevated triose phosphate levels induced by 30 mM glucose in glomeruli from untreated rats were prevented in glomeruli from ARI-fed rats (but not by ARI added in vitro only). *snG3P* levels

in 30 mM glucose-incubated glomeruli were decreased by ARI added in vitro only as well as in glomeruli isolated from ARI-fed rats. Although lactate production by glomeruli (incubated in 30 mM glucose) from ARI-fed rats tended to be lower than that by glomeruli from controls, the difference was not statistically significant.

Discussion

The demonstration that exposure of isolated glomeruli from normal rats to elevated glucose levels in vitro causes a more reduced cytosolic ratio of NADH/NAD^+ is consistent with corresponding effects of elevated glucose levels on cytosolic NADH/NAD^+ in a number of tissues from animals and in human erythrocytes, as noted in the introductory paragraphs of this paper. Furthermore, this glucose-induced glomerular redox change, like that in erythrocytes, granulation tissue, and lens, is prevented or markedly attenuated by aldose reductase inhibitors [6, 18, 25–27, 39, unpublished observations]. It should be noted that glomeruli are comprised of several different cell types (endothelial, epithelial, and mesangial cells) and the metabolic changes observed in the present studies reflect the sum of the changes in all of them. On the other hand, all of these cells are components of glomerular vessels, and the magnitude of the metabolic changes observed supports the likelihood that they reflect the metabolic state in a significant proportion of the glomerular vasculature. These observations have a number of potentially important implications for the pathogenesis of diabetic glomerulopathy.

Metabolic effects of elevated glucose levels

The ~30% increase in lactate production in glomeruli exposed to 30 versus 5 mM glucose is consistent with evidence of glucose-induced increases in lactate production in skin chamber granulation tissue [18] and increased glycolysis in bovine retinal endothelial cells [22], and increased lactate levels in retina and sciatic nerve endoneurium from diabetic rats [40]. Glucose-induced increased lactate production in all of these tissues could be explained by the presence of hexokinase(s) with a relatively high K_m for glucose [41] or release (by elevated glucose levels) of hexokinase inhibition by G6P [42]. The elevated levels of G6P and F6P in 30 mM glucose-incubated glomeruli (Figs. 2 and 4) would be consistent with both possibilities.

The increase in triose phosphate levels induced by 30 mM glucose (Figs. 2, 4, and 8) is consistent with a more reduced cytosolic ratio of NADH/NAD^+ since oxidation of GAP (which is in equilibrium with DHAP and FDP) to 1,3 DPG is coupled to reduction of NAD^+ to NADH (Fig. 1). Since a more reduced cytosolic ratio of NADH/NAD^+ favors *snG3P* synthesis by reduction of DHAP (with NADH serving as the hydrogen donor), the finding that *snG3P* levels were not elevated in 30 mM glucose-incubated glomeruli raises the possibility that utilization of *snG3P* was increased. This interpretation is consistent with evidence that 30 mM glucose increases *de novo* synthesis of DAG in isolated glomeruli [21], since reduction of DHAP to *snG3P* is the first step in one pathway for *de novo* synthesis of DAG (Fig. 1). These observations and interpretations also are consistent with evidence that *snG3P* dehydrogenase activity is not high enough in some tissues (under some conditions) to maintain equilibrium between its substrates (*snG3P* and DHAP) [19, 43].




Fig. 4. Effects of 3 mM pyruvate and 0.07 mM tolrestat on glycolytic intermediates (expressed as nmol/ μ g hydroxyproline, $N = 9$ for each group, mean \pm 1 SD) in glomeruli incubated with 30 mM glucose for one hour at 37°C ($N = 9$ for each group). Significantly different from 5 mM glucose: * $P \leq 0.001$, † $P \leq 0.005$, ‡ $P \leq 0.001$, § $P < 0.05$. Significantly different from 30 mM glucose: †† $P \leq 0.0002$, ††† $P < 0.006$.

The increased sorbitol levels in 30 mM glucose-incubated glomeruli (Fig. 6) is consistent with previous reports [12, 44] and, together with the increased fructose levels (Fig. 6) and the prevention of 30 mM glucose-induced increase in the lactate/pyruvate ratio by aldose reductase inhibitors, supports the hypothesis that increased oxidation of sorbitol to fructose contributes significantly to the more reduced cytosolic ratio of NADH/NAD⁺ in glomeruli exposed to elevated glucose levels.

Effects of aldose reductase inhibitors and pyruvate on glucose-induced metabolic imbalances

Glucose-induced increases in lactate/pyruvate ratios (in glomeruli plus medium) were relatively unaffected by coadministration of ARI in vitro only (Fig. 5), but were prevented in glomeruli from ARI-fed rats (Fig. 7A). These discordant findings may be explained by a relatively slow rate of penetration of glomerular cells by ARI. By the time inhibitory levels were attained in glomeruli exposed to ARI only in vitro, it was too late to normalize the lactate/pyruvate ratio in the medium (most of the lactate and pyruvate are recovered in the medium rather than in the cells). In contrast, excess intracellular sorbitol formed before inhibition of aldose reductase activity was achieved would still be readily oxidized to fructose, that is, "normalized," during the latter part of the incubation. This interpretation is consistent with evidence that sorbitol levels in red cells and other tissues decrease rapidly following normalization of glucose levels [45–47]. The observation that tolrestat decreased tissue lactate/pyruvate ratio in glomeruli incubated in 5 mM glucose suggests that flux of glucose via the sorbitol pathway impacts significantly on glomerular cytosolic NADH/NAD⁺ even at physiological glucose levels.

Aldose reductase utilizes NADPH as the hydrogen donor for the reduction of glucose to sorbitol. This reaction results in a more oxidized ratio of NADP⁺/NADPH (Fig. 1). In contrast, sorbitol dehydrogenase utilizes NAD⁺ as the hydrogen acceptor for oxidation of sorbitol to fructose. This reaction results in a more reduced cytosolic ratio of NADH/NAD⁺ as noted earlier. Therefore, the 30 mM glucose-induced increases in

lactate/pyruvate (Fig. 7) and their prevention by ARI appear to be more closely linked to redox changes resulting from increased oxidation of sorbitol to fructose than to reduction of glucose to sorbitol. This interpretation is supported by several lines of evidence. First, although (as noted earlier) the ratio of pyridine nucleotides in cellular extracts may not accurately reflect the cytosolic ratio of free NADH/NAD⁺, it is noteworthy that 30 mM glucose-induced changes in NADH/NAD⁺ in glomerular extracts are indeed consistent with changes in lactate/pyruvate ratio in the incubation medium (data not shown). Second, topical application of sorbitol to granulation tissue vessels in vivo causes vascular dysfunction, like that induced by glucose, which is prevented by coadministration of pyruvate but not by aldose reductase inhibitors [16, unpublished observations]. Third, the ratio of lactate/pyruvate and the levels of other metabolites (triose phosphates and snG3P) modulated by NADH/NAD⁺ in human erythrocytes are markedly increased during incubation with sorbitol or xylitol (xylitol also is a substrate for sorbitol dehydrogenase) [39, 48], and these increases are prevented by pyruvate [39]. The decrease in snG3P levels (and the trend for triose phosphates to decrease, albeit not statistically significant) in 30 mM glucose-incubated glomeruli from ARI-fed rats (Fig. 8) is consistent with the associated normalization of the cytosolic lactate/pyruvate ratio (Fig. 7B). The observation that snG3P levels in 30 mM glucose-incubated glomeruli from ARI-fed rats were lower than those in 5 and 30 mM glucose-incubated glomeruli from control rats (Fig. 8) may reflect greater inhibition (by ARI) of synthesis than utilization of snG3P (due to differences in oxidation/reduction potentials and equilibrium constants of snG3P dehydrogenase and the enzymes that utilize snG3P for lipid synthesis) [19, 43].

The pyruvate-induced decrease in triose phosphates, snG3P, and sorbitol (Figs. 4 and 6) are consistent with corresponding pyruvate effects on glucose-induced changes in cytosolic NADH/NAD⁺ and on NADH/NAD⁺ modulated metabolite levels (including sorbitol and fructose) in human red cells and in skin chamber granulation tissue [18, 25, 39, unpublished observations]. All of these effects of pyruvate are consistent with




Fig. 5. Effects of 3 mM pyruvate and 0.07 mM tolrestat on lactate, pyruvate, and lactate/pyruvate ratio in glomeruli incubated with 30 mM glucose for one hour at 37°C. Metabolites, expressed as nmol/μg hydroxyproline, are shown for each incubation as the mean ± 1 SD ($N = 9$ for each group). The amount depicted for pyruvate (B) in the 30 mM glucose + 3 mM pyruvate group is an approximate value (indicated by the ~) derived by dividing lactate produced by glomeruli by pyruvate added to the medium. It is shown merely to indicate that addition of 3 mM pyruvate to the medium results in higher pyruvate levels than those produced by glomerular metabolism. Significantly different from 5 mM glucose: * $P < 0.001$, † $P \leq 0.005$.




Fig. 6. Effects of 3 mM pyruvate and 0.07 mM tolrestat on sorbitol, fructose, and myo-inositol accumulation in glomeruli obtained from normal rats after incubation in 30 mM glucose for one hour at 37°C ($N = 6$). Results are expressed as nmol/μg hydroxyproline and are mean ± 1 SD. Significantly different from 5 mM glucose: * $P \leq 0.001$, † $P < 0.014$, ‡ $P < 0.04$. Significantly different from 30 mM glucose: § $P < 0.014$, || $P < 0.011$.

oxidation of NADH to NAD⁺ by pyruvate at a rate faster than NAD⁺ is reduced to NADH by oxidation of sorbitol to fructose (Fig. 1). The apparent failure of pyruvate to increase fructose




Fig. 7. Effects of 5 mM and 30 mM glucose \pm tolrestat on the glomerular tissue lactate/pyruvate ratio (A) or on tissue plus medium lactate/pyruvate ratio (B). Glomeruli used for incubation in 5 or 30 mM glucose + tolrestat were obtained from nondiabetic rats fed tolrestat in their diet (0.02 mmol/kg body weight) for 7 days prior to glomerular isolation. Since these are paired observations, the lactate/pyruvate ratios from each incubation are connected by lines. Mean \pm 1 SD is also indicated.

levels in 30 mM glucose-incubated glomeruli, despite a marked reduction in sorbitol levels, may be due to the presence of fructokinase in the glomeruli and/or in the small number of contaminating tubules, portions of which contain large amounts of the enzyme [49].

These observations raise several important related questions regarding: 1) the reliability of measurements of tissue sorbitol levels alone (without fructose measurements) as an index of sorbitol pathway activity, 2) how much of an increase in sorbitol pathway activity can be tolerated by cells/tissues without pathophysiological consequences, and 3) how readily reversible increases in sorbitol pathway activity contribute to poorly reversible glucose- and diabetes-induced metabolic imbalances, as well as functional and structural angiopathy (in-

cluding glomerulopathy) and neuropathy. Tissue sorbitol levels are determined not only by intracellular glucose levels, but also by the ratio of sorbitol dehydrogenase to aldose reductase enzyme activities [50]. Thus, if the ratio of sorbitol dehydrogenase to aldose reductase in a tissue is high, elevated tissue glucose levels can markedly increase glucose metabolism via the sorbitol pathway to fructose (Fig. 1), and induce sorbitol pathway-mediated cellular injury, with little or no increase in tissue sorbitol content.

The level to which sorbitol pathway activity can be increased without causing metabolic and pathophysiological consequences remains unclear and may vary considerably in different tissues. It appears likely that relatively small increases in sorbitol pathway activity may be sufficient to cause vascular dysfunction in many tissues. This view is supported by the observation that in diabetic rats treated with two different doses of an ARI (AL-1576), both of which normalized sorbitol levels in every tissue examined, only the higher dose decreased microalbuminuria and normalized GFR and vascular albumin permeation in ocular tissues, sciatic nerve, and granulation tissue [5]. These observations have important implications regarding the interpretation of experiments in which ARI have been reported to normalize sorbitol levels but fail to prevent associated glucose-induced changes in protein kinase C activity [20, 23] and diabetes-induced increases in GFR [12] and glomerular structural changes [12]. Differential effects of ARI on metabolic imbalances, functional, and/or structural changes in the same cells and tissues [23, 51–53] may reflect the differential impact of partial normalization of cytosolic NADH/NAD⁺ on dehydrogenase enzymes with different oxidation/reduction potentials and equilibrium constants (and on the relative concentrations of their reduced and oxidized substrates).

The nature of the mechanism(s) by which transient increases in sorbitol pathway activity may induce poorly reversible functional and structural glomerular changes (and other vascular and neural complications of diabetes) also remains unclear. In view of evidence linking increased tissue DAG levels (and associated increased protein kinase C activity) to increased sorbitol pathway activity in granulation tissue [15], it can be postulated that transient elevations in DAG may cause much longer lasting effects through activation and membrane association of protein kinase C and phosphorylation of protein kinase C substrates.

Implications of a more reduced cytosolic NADH/NAD⁺ ratio for the pathogenesis of diabetic nephropathy

One mechanism by which increased sorbitol pathway activity may induce poorly reversible vascular changes is by production of metabolites capable of nonenzymatically glycosylating intracellular constituents such as triose phosphates (which are among the most reactive naturally occurring glycosylating agents in tissues [54]) and fructose 3-phosphate and 3-deoxyglucosone [55]. The likelihood that nonenzymatic glycation may contribute to glucose- and diabetes-induced vascular dysfunction is supported by evidence that aminoguanidine (a nucleophilic hydrazine compound believed to prevent formation of advanced glycation end-products [56]) prevents sorbitol-, glucose-, and diabetes-induced increases in vascular albumin permeation [57–59].

A more reduced cytosolic ratio of NADH/NAD⁺ could impact on at least two important pathways of lipid metabolism




Fig. 8. Glycolytic metabolite levels after incubation in 5 and 30 mM glucose of glomeruli from rats fed ARI in the diet (Methods) prior to glomerular isolation versus metabolite levels in glomeruli isolated from untreated rats with ARI added in vitro only. Metabolite levels are expressed as a ratio of the corresponding level in 5 mM glucose-incubated glomeruli. The dashed line (ratio = 1) represents the metabolite level in 5 mM glucose-incubated glomeruli. Open circles = 30 mM glucose; open triangles = 30 mM glucose + 0.07 mM ARI added to incubation medium in vitro; closed triangles = 30 mM glucose + ARI (tolrestat or AL-1576) added to the diet and also added to the incubation medium (Methods). G6P = glucose 6-phosphate, F6P = fructose 6-phosphate, TP = total triose phosphates (fructose 1,6-bisphosphate, dihydroxyacetone phosphate, and glyceraldehyde 3-phosphate), snG3P = sn-glycerol 3-phosphate, P = pyruvate, L = lactate, and L/P = lactate/pyruvate ratio. G6P and F6P data were not available for the group given ARI in the diet (represented by closed triangles).

Values significantly different from 5 mm glucose at $P \leq$:

○	0.005	0.001	0.025	NS	NS	0.005	0.002
△	0.01	0.005	0.05	0.002	NS	NS	0.002
▲	—	—	NS	0.005	NS	NS	NS
30 mm glucose at $P \leq$:							
△	NS	NS	NS	NS	NS	NS	NS
▲	—	—	NS	0.005	NS	NS	0.05

resulting in: 1) increased *de novo* synthesis of DAG, and 2) impaired β oxidation of long chain fatty acyl CoA in mitochondria. As discussed above, a more reduced cytosolic ratio of NADH/NAD⁺ would favor glucose-induced increased *de novo* synthesis of DAG and associated activation of protein kinase C (Fig. 1) observed in isolated glomeruli exposed to elevated glucose levels in vitro [20, 21]. The putative importance of this cascade of metabolic imbalances is consistent with amelioration by *myo*-inositol of vascular dysfunction (including increases in GFR and in urinary excretion of IgG) in diabetic rats [8, 60], since elevation of tissue *myo*-inositol levels would favor incorporation of DAG into phosphatidylinositol (Fig. 1), thereby lowering DAG levels and normalizing impaired phosphatidylinositol metabolism [61, 62]. This interpretation is consistent with evidence that agonist-induced increases in DAG and CDP-DAG (associated with breakdown of PIP₂) in a variety of tissues are prevented and/or reversed by *myo*-inositol [63–66]. In addition, excess cytosolic reducing equivalents (NADH)

generated by increased oxidation of sorbitol to fructose can be transported into the mitochondria (by the malate-aspartate shuttle). In ischemic myocardium, an increase in mitosomal NADH/NAD⁺ inhibits mitochondrial β oxidation of long-chain fatty acyl CoA (Fig. 1) [11], which accumulate as long-chain acyl esters of CoA and carnitine [67]. These long-chain acyl esters, like DAG, are potent modulators of the activity of several enzymes (including protein kinase C, Na⁺/K⁺-ATPase, and Ca⁺⁺-ATPase) implicated in the pathogenesis of diabetic complications. While the role of this putative cascade of metabolic imbalances in the pathogenesis of diabetic glomerulopathy remains unknown, metabolic imbalances of this nature could explain the apparent links between elevated plasma lipid levels, microalbuminuria, and nephropathy in diabetic humans and animals [68–71].

In conclusion, these studies demonstrate that elevated glucose levels lead to a more reduced ratio of cytosolic NADH/NAD⁺ in glomeruli which is linked to increased flux of glucose

via the sorbitol pathway. They also suggest several candidate mechanisms by which a more reduced glomerular cytosolic ratio of NADH/NAD⁺ may modulate lipid metabolism and nonenzymatic glycation and contribute to the pathogenesis of diabetic nephropathy. While these observations implicate increased sorbitol pathway metabolism in the pathogenesis of glomerular dysfunction in diabetic animals, further studies are needed to elucidate the role of sorbitol pathway metabolism in the pathogenesis of diabetic glomerulopathy in human subjects.

Acknowledgments

This research was supported by grants from the National Institutes of Health (EY06600, HL39934, DK20579) and by the Kilo Diabetes and Vascular Research Foundation. Portions of this research were supported by Pfizer Central Research and by Eli Lilly and Company. We thank Dr. M.A. Province for assistance with the statistical analysis of the data, and Wyeth-Ayerst and Alcon for providing the aldose reductase inhibitors used in these studies. We acknowledge the secretarial assistance of Ms. A. Stone and the photographic assistance of Mr. J. Fulford.

Reprint requests to Ronald G. Tilton, Ph.D., Washington University School of Medicine, Department of Pathology - Box 8118, 660 South Euclid Avenue, St. Louis, Missouri 63110, USA.

Appendix. List of abbreviations

ARI	aldose reductase inhibitor
CDP-DAG	cytidine diphosphate-DAG
DAG	1,2-diacyl- <i>sn</i> -glycerol
DHAP	dihydroxyacetone phosphate
1,3 DPG	1,3-bisphosphoglycerate
FDP	fructose 1,6-bisphosphate
F6P	fructose 6-phosphate
GAP	glyceraldehyde 3-phosphate
G6P	glucose 6-phosphate
KH	Krebs-HEPES buffer
L	lactate
L/P	lactate/pyruvate ratio
P	pyruvate
PIP ₂	phosphatidylinositol-bisphosphate
<i>sn</i> G3P	<i>sn</i> -glycerol 3-phosphate
TP	total triose phosphates

References

- MOGENSEN CE: Glomerular filtration rate and renal plasma flow in short-term and long-term juvenile diabetes mellitus. *Scand J Clin Lab Invest* 28:91-100, 1971
- PARVING H-H, VIBERTI GC, KEEN H, CHRISTIANSEN JS, LASSEN NA: Hemodynamic factors in the genesis of diabetic microangiopathy. *Metabolism* 32:943-949, 1983
- MOGENSEN CE: Early glomerular hyperfiltration in insulin-dependent diabetics and late nephropathy. *Scand J Clin Lab Invest* 46:201-206, 1986
- ZATZ R, BRENNER BM: Pathogenesis of diabetic microangiopathy: The hemodynamic view. *Am J Med* 80:443-453, 1987
- TILTON RG, CHANG K, PUGLIESE G, EADES DM, PROVINCE MA, SHERMAN WR, KILO C, WILLIAMSON JR: Prevention of hemodynamic and vascular albumin filtration changes in diabetic rats by aldose reductase inhibitors. *Diabetes* 38:1258-1270, 1989
- WILLIAMSON JR, KILO C, TILTON RG: Mechanisms of glucose- and diabetes-induced vascular dysfunction, in *Hyperglycemia, Diabetes and the Vascular Wall*, edited by RUDERMAN NB, BROWNLEE MA, WILLIAMSON JR, New York, Oxford University Press, 1991
- BEYER-MEARS A, CRUZ E, EDELST T, VARAGIANNIS E: Diminished proteinuria in diabetes mellitus by sorbinil, an aldose reductase inhibitor. *Pharmacology* 32:52-60, 1980
- GOLDFARB S, SIMMONS DA, KERN K: Amelioration of glomerular hyperfiltration in acute experimental diabetes by dietary *myo*-inositol and by an aldose reductase inhibitor. *Trans Assoc Am Physiol* 99:67-72, 1986
- BANK N, MOWER P, AYNEDJIAN HS, WILKES BM, SILVERMAN S: Sorbinil prevents glomerular hyperperfusion in diabetic rats. *Am J Physiol* 256:F1000-F1006, 1989
- PUGLIESE G, TILTON RG, SPEEDY A, CHANG K, PROVINCE MA, KILO C, WILLIAMSON JR: Vascular filtration function in galactose-fed versus diabetic rats: The role of polyol pathway activity. *Metabolism* 39:690-697, 1990
- PUGLIESE G, TILTON RG, WILLIAMSON JR: Glucose-induced metabolic imbalances in the pathogenesis of diabetic vascular disease. *Diab/Metabol Rev* 7:35-59, 1991
- DANIELS BS, HOSTETTER TH: Aldose reductase inhibition and glomerular abnormalities in diabetic rats. *Diabetes* 38:981-986, 1989
- PEDERSEN MM, CHRISTIANSEN JS, MOGENSEN CK: Reduction of glomerular hyperfiltration in normoalbuminuric IDDM patients by 6 mo of aldose reductase inhibition. *Diabetes* 40:527-531, 1991
- WILLIAMSON JR, OSTROW E, EADES DM, CHANG K, ALLISON W, KILO C, SHERMAN WR: Glucose-induced microvascular functional changes in nondiabetic rats are stereospecific and are prevented by an aldose reductase inhibitor. *J Clin Invest* 85:1167-1172, 1990
- WOLF BA, WILLIAMSON JR, EASOM RA, CHANG K, SHERMAN WR, TURK J: Diacylglycerol accumulation and microvascular abnormalities induced by elevated glucose levels. *J Clin Invest* 87:31-38, 1990
- WILLIAMSON JR, CHANG K, OSTROW E, ALLISON W, HARLOW J, KILO C: Sorbitol-induced increases in vascular albumin clearance (VAC) are prevented by pyruvate but not *myo*-inositol. (abstract) *Diabetes* 38:94A, 1989
- WILLIAMSON JR, CHANG K, OSTROW E, TIMM V, ALLISON W: Normalization of D-glucose-induced increases in blood flow (BF) and ¹²⁵I-albumin permeation (IAP) by *myo*-inositol (MI) and by pyruvate. (abstract) *Diabetes* 37:95A, 1988
- KAWAMURA T, SMITH S, WILLIAMSON JR: Glucose-induced metabolic changes in tissue chamber granulation tissue. (abstract) *Diabetes* 39:192A, 1990
- WILLIAMSON DH, LUND P, KREBS HA: The redox state of free nicotinamide-adenine dinucleotide in the cytoplasm and mitochondria of rat liver. *Biochem J* 103:514-527, 1967
- CRAVEN PA, DERUBERTIS FR: Protein kinase C is activated in glomeruli from streptozotocin diabetic rats: Possible mediation by glucose. *J Clin Invest* 83:1667-1675, 1989
- CRAVEN PA, DAVIDSON CM, DERUBERTIS FR: Increase in diacylglycerol mass in isolated glomeruli by glucose from *de novo* synthesis of glycerolipids. *Diabetes* 39:667-674, 1990
- LEE T-S, SALTSMAN KA, OHASHI H, KING GL: Activation of protein kinase C by elevation of glucose concentration: Proposal for a mechanism in the development of diabetic vascular complications. *Proc Natl Acad Sci USA* 86:5141-5145, 1989
- LEE T-S, MACGREGOR LC, FLUHARTY SJ, KING GL: Differential regulation of protein kinase C and (Na,K)-adenosine triphosphatase activities by elevated glucose levels in retinal capillary endothelial cells. *J Clin Invest* 83:90-94, 1989
- OKUMURA K, AKIYAMA N, HASHIMOTO H, OGAWA K, SATAKE T: Alteration of 1,2-diacylglycerol content in myocardium from diabetic rats. *Diabetes* 37:1168-1172, 1988
- TRAVIS SF, MORRISON AD, CLEMENTS RS JR, WINEGRAD AI, OSKI FA: Metabolic alterations in the human erythrocyte produced by increases in glucose concentration: The role of the polyol pathway. *J Clin Invest* 50:2104-2112, 1971
- KINOSHITA JH, FUKUSHI S, KADOR P, MEROLA LO: Aldose reductase in diabetic complications of the eye. *Metabolism* 28:462-469, 1979
- CHENG H-M, GONZALEZ RG, VON SALTZA I, CHYLACK LT JR, HUTSON NJ: Glucose flux and the redox state of pyridine dinucleotides in the rat lens. *Exp Eye Res* 46:947-952, 1988

28. CHRISTIANSEN JS, FRANDSEN M, PARVING H-H: Effect of intravenous glucose infusion on renal function in normal man and in insulin-dependent diabetics. *Diabetologia* 21:368-373, 1981
29. WOODS LL, MIZELLE HL, HALL JE: Control of renal hemodynamics in hyperglycemia: Possible role of tubuloglomerular feedback. *Am J Physiol* 252:F65-F73, 1987
30. TREVISAN R, NOSADINI R, FIORETTO P, VELUSSI M, AVOGARO A, DUNER E, LORI E, DORIA A, MERKEL C, VALERIO A, CREPALDI G: Metabolic control of kidney hemodynamics in normal and insulin-dependent diabetic subjects: Effects of acetoacetic, lactic, and acetic acids. *Diabetes* 36:1073-1081, 1987
31. SCHLONDORFF D, ROCZNIK S, SATRIANO JA, FOLKERT VW: Prostaglandin synthesis by isolated rat glomeruli: Effect of angiotensin II. *Am J Physiol* 238:F486-F495, 1980
32. BERGMAYER H-U: *Methods of Enzymatic Analysis*. Weinheim, Verlag Chemie, 1962
33. LOWRY OH, PASSONEAU JV: *A Flexible System of Enzymatic Analysis*. Orlando, Academic Press, 1972
34. KIVIRIKKO KI, LAITINEN O, PROCKOP DJ: Modifications for a specific assay of hydroxypyroline in urine. *Anal Biochem* 19:249-255, 1967
35. CHI MM-Y, PUSATERI ME, CARTER JG, NORRIS BJ, MCDUGAL DB JR, LOWRY OH: Enzymatic assays for 2-deoxyglucose and 2-deoxyglucose 6-phosphate. *Anal Biochem* 161:508-513, 1987
36. KREBS HA: The redox state of nicotinamide adenine dinucleotide in the cytoplasm and mitochondria of rat liver. *Adv Enzymol Reg* 5:409-434, 1967
37. EADES DM, WILLIAMSON JR, SHERMAN WR: Rapid analysis of sorbitol, galactitol, mannitol, and myo-inositol mixtures from biological sources. *J Chromatogr* 490:1-8, 1989
38. ANDERSON TW: *An Introduction to Multivariate Statistical Analysis*. New York, Wiley, 1958
39. FRANGOS M, SMITH S, SANTIAGO J, KILO C: Sorbitol-induced imbalances in glycolysis in human erythrocytes are reduced by pyruvate. (abstract) *Diabetes* 39 (Suppl. 1):274A, 1990
40. POULSON R, MIRRELES DJ, EARL DCN, HEALTH H: The effects of an aldose reductase inhibitor upon the sorbitol pathway, fructose-1-phosphate and lactate in the retina and nerve of streptozotocin-diabetic rats. *Exp Eye Res* 36:751-760, 1983
41. GROSSBARD L, SCHIMKE RT: Multiple hexokinases of rat tissues: Purification and comparison of soluble forms. *J Biol Chem* 241:3546-3560, 1966
42. FUJII S, BEUTLER E: High glucose concentrations partially release hexokinase from inhibition by glucose 6-phosphate. *Proc Natl Acad Sci USA* 82:1552-1554, 1985
43. WILLIAMSON JR, KREISBERG RA, FELTS PW: Mechanism for the stimulation of gluconeogenesis by fatty acids in perfused rat liver. *Proc Natl Acad Sci USA* 56:247-254, 1966
44. BEYER-MEARS A, KU L, COHEN MP: Glomerular polyol accumulation in diabetes and its prevention by oral sorbinil. *Diabetes* 33:604-607, 1984
45. NAGASAKA Y, FUJII S, KANEKO T: Human erythrocyte sorbitol metabolism and the role of sorbitol dehydrogenase. *Diabetologia* 31:766-770, 1988
46. ROGERS D, ROWOLD E, CHANG K, TOMLINSON M, SHERMAN WR, GAVIN J, KILO C, WILLIAMS JR: Effect of rapid normalization of plasma glucose levels on microvascular dysfunction and polyol metabolism in diabetic rats. *Diabetes* 37:1689-1694, 1988
47. PUGLIESE G, TILTON RG, CHANG K, SPEEDY A, PROVINCE M, EADES DM, LACY PE, KILO C, WILLIAMSON JR: Effects of islet isografts on hemodynamic and vascular filtration changes in diabetic rats. *Diabetes* 39:323-332, 1990
48. ASAKURA T, ADACHI K, MINAKAMI S, YOSHIKAWA H: Non-glycolytic sugar metabolism in human erythrocytes. I. Xylitol metabolism. *J Biochem (Tokyo)* 62:184-193, 1967
49. BURCH HB, CHOI S, DENCE CN, ALVEY TR, COLE BR, LOWRY OH: Metabolic effects of large fructose loads in different parts of the rat nephron. *J Biol Chem* 255:8239-8244, 1980
50. GRIFFIN BW, MCNATT LG, YORK BM JR: Characterization of aldose reductase activities from human and animal sources by a sensitive fluorescence assay, in *Enzymology and Molecular Biology and Carbonyl Metabolism*, edited by WEINER H, New York, Alan R. Liss, Inc., 1987, pp. 325-340
51. YEH L-A, RAFFORD CE, GODDU KJ, ASHTON MA, BEYER TA, HUTSON NJ: Na⁺-K⁺-ATPase pumping activity is not directly linked to myo-inositol levels after sorbinil treatment in lenses of diabetic rats. *Diabetes* 36:1414-1419, 1987
52. MCCAULEY ML, SREDY J, MILLEN J, ACKERMAN DM, DVORNIK D: Prevention of urinary albumin excretion in 6 month streptozotocin-diabetic rats with the aldose reductase inhibitor tolrestat. *J Diabetic Compl* 2:16-18, 1988
53. TILTON RG, PUGLIESE G, WILLIAMSON JR: Diabetes-induced glomerular structural changes in rats are not prevented by sorbinil. (abstract) *Diabetes* 38:94A, 1989
54. STEVENS VJ, VLASSARA H, ABATI A, CERAMI A: Nonenzymatic glycosylation of hemoglobin. *J Biol Chem* 252:2998-3002, 1977
55. SZWARGOLD BS, KAPPLER F, BROWN TR: Identification of fructose 3-phosphate in the lens of diabetic rats. *Science* 247:451-454, 1990
56. BROWNEE M, CERAMI A, VLASSARA H: Advanced glycosylation end products in tissue and the biochemical basis of diabetic complications. *N Engl J Med* 318:1315-1321, 1988
57. IDO Y, CHANG K, OSTROW E, ALLISON W, KILO C, TILTON RG: Aminoguanidine prevents regional blood flow increases in streptozotocin-diabetic rats. (abstract) *Diabetes* 39:93A, 1990
58. TILTON RG, CHANG K, OSTROW E, ALLISON W, WILLIAMSON JR: Aminoguanidine reduces increased ¹³¹I-albumin permeation of retinal and uveal vessels in streptozotocin-diabetic rats. (abstract) *Invest Ophthalmol Vis Sci* 31:342, 1990
59. CHANG K, ALLISON W, HARLOW J, WILLIAMSON JR: Aminoguanidine prevents glucose- and sorbitol-induced vascular dysfunction in skin chamber granulation tissue. (abstract) in *Proceedings of 14th Congress of the International Diabetes Federation*, Washington, DC, 1991
60. PUGLIESE G, TILTON RG, SPEEDY A, SANTARELLI E, EADES DM, PROVINCE MA, KILO C, SHERMAN WR, WILLIAMSON JR: Modulation of hemodynamic and vascular filtration changes in diabetic rats by dietary myo-inositol. *Diabetes* 39:312-322, 1990
61. WINEGRAD AI: Does a common mechanism induce the diverse complications of diabetes? *Diabetes* 36:396-406, 1987
62. SIMMONS DA, WINEGRAD AI: Mechanism of glucose-induced (Na⁺, K⁺)-ATPase inhibition in aortic wall of rabbits. *Diabetologia* 32:402-408, 1989
63. DRUMMOND AH, RAEBURN CA: The interaction of lithium with thyrotropin-releasing hormone-stimulated lipid metabolism in GH3 pituitary tumour cells. *Biochem J* 224:129-136, 1984
64. DOWNES CP, STONE MA: Lithium-induced reduction in intracellular inositol supply in cholinergically stimulated parotid gland. *Biochem J* 234:199-204, 1986
65. BERRIDGE MJ, DOWNES CP, HANLEY MJ: Neural and developmental actions of lithium: A unifying hypothesis. *Cell* 59:411-419, 1989
66. GODFREY PP: Potentiation by lithium of CMP-phosphatidate formation in carbachol-stimulated rat cerebral-cortical slices and its reversal by myo-inositol. *Biochem J* 258:621-624, 1989
67. CORR PB, GROSS RW, SOBEL BE: Amphipathic metabolites and membrane dysfunction in ischemic myocardium. *Circ Res* 55:135-154, 1984
68. KEANE WF, KASISKE BL, O'DONNELL MP: Hyperlipidemia and the progression of renal disease. *Am J Clin Nutr* 47:157-160, 1988
69. COLLINS VR, DOWSE GK, FINCH CF, ZIMMET PZ, LINNANE AW: Prevalence and risk factors for micro- and macroalbuminuria in diabetic subjects and entire population of Nauru. *Diabetes* 38:1602-1610, 1989
70. DUNN FL: Hyperlipidemia in diabetes mellitus. *Diab/Metab Rev* 6:47-61, 1990
71. TUNG P, GINIER P, LEVIN SR, HERSHMAN JD, HERSHMAN JM: Clinical characteristics associated with microalbuminuria in an adult diabetic population. *J Diabetes Compl* 4:15-20, 1990